

Product datasheet for TA367069S

HCAR2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human HCAR2Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: hydroxycarboxylic acid receptor 2

Database Link: Entrez Gene 338442 Human

Q8TDS4

Background: Acts as a high affinity receptor for both nicotinic acid (also known as niacin) and (D)-beta-

hydroxybutyrate and mediates increased adiponectin secretion and decreased lipolysis through G(i)-protein-mediated inhibition of adenylyl cyclase. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet. Mediates nicotinic acid-induced apoptosis in mature neutrophils. Receptor activation by nicotinic acid results in reduced cAMP levels which may affect activity of cAMP-dependent protein kinase A and phosphorylation of target proteins, leading to neutrophil apoptosis. The rank order of potency for the displacement of nicotinic acid binding is 5-methyl pyrazole-3-carboxylic acid = pyridine-3-acetic acid > acifran > 5-methyl nicotinic acid = acipimox >>

nicotinuric acid = nicotinamide.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

GPR109A; HCA2; HM74a; HM74b; NIACR1; Puma-g; PUMAG

Product images:




Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA367069] (HCAR2 Antibody) at dilution 1/50 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA367069] (HCAR2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA367069] (HCAR2 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA367069] (HCAR2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)