

Product datasheet for TA367071

GPD2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 20-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human GPD2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: glycerol-3-phosphate dehydrogenase 2

Database Link: Entrez Gene 2820 Human

P43304

Background: The protein encoded by this gene localizes to the inner mitochondrial membrane and

catalyzes the conversion of glycerol-3-phosphate to dihydroxyacetone phosphate, using FAD as a cofactor. Along with GDP1, the encoded protein constitutes the glycerol phosphate shuttle, which reoxidizes NADH formed during glycolysis. Two transcript variants encoding

the same protein have been found for this gene.

Synonyms: AA408484; Al448216; AU021455; AW494132; Gdm1; Gpd-m; GPDH; Gpdh-m; mtGPDH; TISP38


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA367071 (GPD2 Antibody) at dilution 1/20 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA367071 (GPD2 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA367071 (GPD2 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA367071 (GPD2 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)