

# **Product datasheet for TA367101**

## Importin 7 (IPO7) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela cells

IHC: 100-300

Positive control: Human gasrtic cancer

Predicted cell location: Cytoplasm or Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human IPO7

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 120 kDa
Gene Name: importin 7

Database Link: Entrez Gene 10527 Human

<u>095373</u>

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

The importin-alpha/beta complex and the GTPase Ran mediate nuclear import of proteins with a classical nuclear localization signal. The protein encoded by this gene is a member of a class of approximately 20 potential Ran targets that share a sequence motif related to the Ran-binding site of importin-beta. Similar to importin-beta, this protein prevents the activation of Ran's GTPase by RanGAP1 and inhibits nucleotide exchange on RanGTP, and also binds directly to nuclear pore complexes where it competes for binding sites with importin-beta and transportin. This protein has a Ran-dependent transport cycle and it can cross the nuclear envelope rapidly and in both directions. At least four importin beta-like transport receptors, namely importin beta itself, transportin, RanBP5 and RanBP7, directly bind and import ribosomal proteins.

Synonyms: FLJ14581; Imp7; MGC138673; RANBP7

### **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 μg Lane: Hela cells


Primary antibody: TA367101 (IPO7 Antibody) at

dilution 1/300

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 minutes




Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using TA367101 (IPO7 Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using TA367101 (IPO7 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA367101 (IPO7 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA367101 (IPO7 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)