

Product datasheet for TA367403S

ATF1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela cell lysate

IHC: 25-100

Positive control: Human lung cancer Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human ATF1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 29 kDa

Gene Name: activating transcription factor 1

Database Link: Entrez Gene 466 Human

P18846

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

This gene encodes an activating transcription factor, which belongs to the ATF subfamily and bZIP (basic-region leucine zipper) family. It influences cellular physiologic processes by regulating the expression of downstream target genes, which are related to growth, survival, and other cellular activities. This protein is phosphorylated at serine 63 in its kinase-inducible domain by serine/threonine kinases, cAMP-dependent protein kinase A, calmodulin-dependent protein kinase I/II, mitogen- and stress-activated protein kinase and cyclin-dependent kinase 3 (cdk-3). Its phosphorylation enhances its transactivation and transcriptional activities, and enhances cell transformation. Fusion of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates chimeric proteins in angiomatoid fibrous histiocytoma and clear cell sarcoma. This gene has a pseudogene on chromosome 6.

Synonyms: EWS-ATF1; FUS/ATF-1; TREB36

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane: Hela cell lysate


Primary antibody: [TA367403] (ATF1 Antibody) at

dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA367403] (ATF1 Antibody) at dilution 1/35 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA367403] (ATF1 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA367403] (ATF1 Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA367403] (ATF1 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)