

Product datasheet for TA367419

COX4 (COX4I1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: NIH/3T3, Hela and 293T cell lysates

IHC: 20-100

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human COX4I1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 20 kDa

Gene Name: cytochrome c oxidase subunit 4l1

Database Link: Entrez Gene 1327 Human

P13073



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer and proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit IV isoform 1 of the human mitochondrial respiratory chain enzyme. It is located at the 3' of the NOC4 (neighbor of COX4) gene in a head-to-head orientation, and shares a promoter with it.

Synonyms: COX4; COXIV; MGC72016

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-3: NIH/3T3 Hela and 293T cell lysates


Primary antibody: TA367419 (COX4I1 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 3 seconds




Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA367419 (COX4I1 Antibody) at dilution 1/20 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA367419 (COX4I1 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA367419 (COX4I1 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA367419 (COX4I1 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)