

## Product datasheet for **TA367475S**

### **Bim (BCL2L11) Rabbit Polyclonal Antibody**

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: K562, A431, HeLa, A549, Raji cell, Rat kidney tissue lysates IHC: 20-100 Positive control: Human ovarian cancer Predicted cell location: Cytoplasm
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human BCL2L11
Formulation:	pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	22 kDa
Gene Name:	BCL2 like 11
Database Link:	<a href="#">Entrez Gene 10018 Human O43521</a>




[View online »](#)

**Background:**


The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The protein encoded by this gene contains a Bcl-2 homology domain 3 (BH3). It has been shown to interact with other members of the BCL-2 protein family and to act as an apoptotic activator. The expression of this gene can be induced by nerve growth factor (NGF), as well as by the forkhead transcription factor FKHR-L1, which suggests a role of this gene in neuronal and lymphocyte apoptosis. Transgenic studies of the mouse counterpart suggested that this gene functions as an essential initiator of apoptosis in thymocyte-negative selection. Several alternatively spliced transcript variants of this gene have been identified.

**Synonyms:**


BAM; Bcl2-L-11; BIM; BIM-alpha6; BIM-beta6; BIM-beta7; BimEL; BimL; BOD; OTTHUMP00000197013

**Product images:**



Gel: 12%SDS-PAGE  
 Lysate: 40 µg  
 Lane 1-6: K562  
 A431  
 HeLa  
 A549  
 Raji cell  
 Rat kidney tissue lysates  
 Primary antibody: [TA367475] (BCL2L11 Antibody) at dilution 1/500  
 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution  
 Exposure time: 3 minutes




Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA367475] (BCL2L11 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA367475] (BCL2L11 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA367475] (BCL2L11 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA367475] (BCL2L11 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)