

# **Product datasheet for TA367829**

## Her2 (ERBB2) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Hela and HUVEC cell lysates

IHC: 20-100

Positive control: Human lung cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human ERBB2

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 138 kDa

**Gene Name:** erb-b2 receptor tyrosine kinase 2

Database Link: Entrez Gene 2064 Human

P04626

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

The ErbB2 (HER2) proto-oncogene encodes a 185 kDa transmembrane, receptor-like glycoprotein with intrinsic tyrosine kinase activity. While ErbB2 lacks an identified ligand, ErbB2 kinase activity can be activated in the absence of a ligand when overexpressed and through heteromeric associations with other ErbB family members. Amplification of the ErbB2 gene and overexpression of its product are detected in almost 40% of human breast cancers. Binding of the c-Cbl ubiquitin ligase to ErbB2 at Tyr1112 leads to ErbB2 poly-ubiquitination and enhances degradation of this kinase. ErbB2 is a key therapeutic target in the treatment of breast cancer and other carcinomas and targeting the regulation of ErbB2 degradation by the c-Cbl-regulated proteolytic pathway is one potential therapeutic strategy. Phosphorylation of the kinase domain residue Tyr877 of ErbB2 (homologous to Tyr416 of pp60c-Src) may be involved in regulating ErbB2 biological activity. The major autophosphorylation sites in ErbB2 are Tyr1248 and Tyr1221/1222; phosphorylation of these sites couples ErbB2 to the Ras-Raf-MAP kinase signal transduction pathway.

Synonyms:

C-erbB-2; CD340; erbB-2; HER-2; HER-2/neu; HER2; herstatin; NEU; NGL; p185erbB2; TKR1


### **Product images:**




Gel: 6%SDS-PAGE Lysate: 40 µg Lane 1-2: Hela and HUVEC cell lysates Primary antibody: TA367829 (ERBB2 Antibody) at dilution 1/350 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 1 minute






Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA367829 (ERBB2 Antibody) at dilution 1/35 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA367829 (ERBB2 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA367829 (ERBB2 Antibody) at dilution 1/35 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA367829 (ERBB2 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)