

Product datasheet for TA368060

MBD2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human cerebrum tissue and Mouse brain tissue lysates

IHC: 30-150

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human MBD2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 43 kDa

Gene Name: methyl-CpG binding domain protein 2

Database Link: Entrez Gene 8932 Human

Q9UBB5



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of binding specifically to methylated DNA. MECP2, MBD1 and MBD2 can also repress transcription from methylated gene promoters. The protein encoded by this gene may function as a mediator of the biological consequences of the methylation signal. It is also reported that the this protein functions as a demethylase to activate transcription, as DNA methylation causes gene silencing. Two transcript variants encoding different isoforms have been found for this gene.

Synonyms:

Demethylase; DKFZp586O0821; DMTase; NY-CO-41

Product images:




Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-2: Human cerebrum tissue and Mouse
brain tissue lysates
Primary antibody: TA368060 (MBD2 Antibody) at
dilution 1/300
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution


Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA368060 (MBD2 Antibody) at dilution 1/30 (Original magnification: ×200)

Exposure time: 10 minutes






Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA368060 (MBD2 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA368060 (MBD2 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA368060 (MBD2 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)