

Product datasheet for TA368069

RPL14 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: 293T Hela 231 HEPG2 and A431 cell lysates

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human RPL14

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 23 kDa

Gene Name: ribosomal protein L14

Database Link: Entrez Gene 9045 Human

P50914



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L14E family of ribosomal proteins. It contains a basic region-leucine zipper (bZIP)-like domain. The protein is located in the cytoplasm. This gene contains a trinucleotide (GCT) repeat tract whose length is highly polymorphic; these triplet repeats result in a stretch of alanine residues in the encoded protein. Transcript variants utilizing alternative polyA signals and alternative 5'-terminal exons exist but all encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Synonyms:

CAG-ISL-7; CTG-B33; hRL14; L14; MGC88594; RL14

Product images:



Gel: 12%SDS-PAGE Lysate: 40 μg Lane 1-5: 293T Hela

231

HEPG2 and A431 cell lysates

Primary antibody: TA368069 (RPL14 Antibody) at

dilution 1/800

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 seconds