

## **Product datasheet for TA368115S**

## Presenilin 1 (PSEN1) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: K562 cell, Mouse brain tissue lysates

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human PSEN1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

**Stability:** 1 year **Predicted Protein Size:** 53 kDa

**Gene Name:** presenilin 1

Database Link: Entrez Gene 5663 Human

P49768

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






## Background:

Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1; PSEN2) or in the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the longer form of amyloid-beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes. Several alternatively spliced transcript variants encoding different isoforms have been identified for this gene, the full-length nature of only some have been determined.


Synonyms:

AD3; FAD; OTTHUMP00000180589; PS-1; PS1; PSNL1; S182

## **Product images:**




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: K562 cell Mouse brain tissue lysates Primary antibody: [TA368115] (PSEN1 Antibody) at dilution 1/800 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution Exposure time: 3 minutes



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368115] (PSEN1 Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368115] (PSEN1 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)