

## **Product datasheet for TA368133S**

# **Lipin 1 (LPIN1) Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: NIH/3T3 IK562 and HepG2 cell lysates

IHC: 50-300

Positive control: Human gastric cancer Predicted cell location: Nucleus membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human LPIN1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol


**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 99 kDa
Gene Name: lipin 1

Database Link: Entrez Gene 23175 Human

Q14693



OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

This gene encodes a magnesium-ion-dependent phosphatidic acid phosphohydrolase enzyme that catalyzes the penultimate step in triglyceride synthesis including the dephosphorylation of phosphatidic acid to yield diacylglycerol. Expression of this gene is required for adipocyte differentiation and it also functions as a nuclear transcriptional coactivator with some peroxisome proliferator-activated receptors to modulate expression of other genes involved in lipid metabolism. Mutations in this gene are associated with metabolic syndrome, type 2 diabetes, and autosomal recessive acute recurrent myoglobinuria (ARARM). This gene is also a candidate for several human lipodystrophy syndromes. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice variants have been described but their full-length structures have not been determined.

Synonyms: DKFZp781P1796; KIAA0188; PAP1

### **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 µg Lane 1-3: NIH/3T3 K562 and HepG2 cell lysates

Primary antibody: [TA368133] (LPIN1 Antibody) at

dilution 1/850


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 seconds

Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA368133] (LPIN1 Antibody) at dilution 1/60 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA368133] (LPIN1 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)