

Product datasheet for TA368168S

ALX4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 20-100

Positive control: Human liver cancer Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human ALX4

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: ALX homeobox 4

Database Link: <u>Entrez Gene 60529 Human</u>

Q9H161

Background: This gene encodes a paired-like homeodomain transcription factor expressed in the

mesenchyme of developing bones, limbs, hair, teeth, and mammary tissue. Mutations in this gene cause parietal foramina 2 (PFM2); an autosomal dominant disease characterized by deficient ossification of the parietal bones. Mutations in this gene also cause a form of frontonasal dysplasia with alopecia and hypogonadism; suggesting a role for this gene in craniofacial development, mesenchymal-epithelial communication, and hair follicle

development. Deletion of a segment of chromosome 11 containing this gene, del(11)(p11p12), causes Potocki-Shaffer syndrome (PSS); a syndrome characterized by craniofacial anomalies, mental retardation, multiple exostoses, and genital abnormalities in males. In mouse, this gene has been shown to use dual translation initiation sites located 16 codons apart.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: FPP; KIAA1788; PFM; PFM2

Product images:



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368168] (ALX4 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368168] (ALX4 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)