

# **Product datasheet for TA368171S**

#### Troduct datasticct for TASOOTA is

## **H2BC1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: TM4, A549 and PC-3 cell, Human prostate tissue and Mouse adipose

tissue lysates IHC: 30-150

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

**Reactivity:** Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide of human H2BC1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 14 kDa

**Gene Name:** histone cluster 1, H2ba

**Database Link:** Entrez Gene 255626 Human

Q96A08



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a testis/sperm-specific member of the histone H2B family. Transcripts from this gene contain a palindromic termination element. [provided by RefSeq, Aug 2015]

Synonyms:

bA317E16.3; H2BFU; STBP; TSH2B

## **Product images:**




Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-5: TM4 A549 and PC-3 cell Human prostate tissue an

Human prostate tissue and Mouse adipose tissue lysates

Primary antibody: [TA368171] (H2BC1 Antibody) at dilution 1/300


Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 3 seconds




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA368171] (H2BC1 Antibody) at dilution 1/25 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA368171] (H2BC1 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368171] (H2BC1 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368171] (H2BC1 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)