

## **Product datasheet for TA368197S**

## **DOC2B Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 20-100

Positive control: Human liver cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human DOC2BFormulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: double C2 domain beta

Database Link: Entrez Gene 8447 Human

Q14184

Background: There are at least two protein isoforms of the Double C2 protein, namely alpha (DOC2A) and

beta (DOC2B), which contain two C2-like domains. DOC2A and DOC2B are encoded by different genes; these genes are at times confused with the unrelated DAB2 gene which was initially named DOC-2. DOC2B is expressed ubiquitously and is suggested to be involved in

Ca(2+)-dependent intracellular vesicle trafficking in various types of cells.

Synonyms: Doc2-beta; DOC2BL




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368197] (DOC2B Antibody) at dilution 1/20 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368197] (DOC2B Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA368197] (DOC2B Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA368197] (DOC2B Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)