

Product datasheet for **TA368243**

ENT2 (SLC29A2) Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: MCF-7 cell lysate IHC: 25-50 Positive control: Human gastric cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human SLC29A2
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	50 kDa
Gene Name:	solute carrier family 29 member 2
Database Link:	Entrez Gene 3177 Human Q14542
Background:	The uptake of nucleosides by transporters, such as SLC29A2, is essential for nucleotide synthesis by salvage pathways in cells that lack de novo biosynthetic pathways. Nucleoside transport also plays a key role in the regulation of many physiologic processes through its effect on adenosine concentration at the cell surface.
Synonyms:	DER12; ENT2; HNP36




[View online »](#)


Product images:




Gel: 8%SDS-PAGE
Lysate: 40 μ g
Lane: MCF-7 cell lysate
Primary antibody: TA368243 (SLC29A2 Antibody) at dilution 1/400
Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution
Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA368243 (SLC29A2 Antibody) at dilution 1/25 (Original magnification: \times 200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA368243 (SLC29A2 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA368243 (SLC29A2 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA368243 (SLC29A2 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)