

Product datasheet for TA368265S

HLAB (HLA-B) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: A549 cell lysate

IHC: 30-150

Positive control: Human cervical cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human HLA-B

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 40 kDa

Gene Name: major histocompatibility complex, class I, B

Database Link: Entrez Gene 3106 Human

P01889



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

HLA-B belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon 1 encodes the leader peptide, exon 2 and 3 encode the alpha1 and alpha2 domains, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Hundreds of HLA-B alleles have been described.

Synonyms: AS; HLAB; SPDA1

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: A549 cell lysate


Primary antibody: [TA368265] (HLA-B Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA368265] (HLA-B Antibody) at dilution 1/30 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA368265] (HLA-B Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)