

Product datasheet for TA368410S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

ATP5ME Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human heart tissue, Human fetal liver tissue, 293T cell, PC-3 cell, Human

liver tissue lysates

IHC: 50-300

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human ATP5MEFormulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 8 kDa

Gene Name: ATP synthase, H+ transporting, mitochondrial Fo complex subunit E

Database Link: Entrez Gene 521 Human

P56385






Background:

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the e subunit of the Fo complex. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Jun 2010]

Synonyms: ATP5K; MGC12532


Product images:




Gel: 12%SDS-PAGE
Lysate: 40 µg
Lane 1-5: Human heart tissue
Human fetal liver tissue
293T cell
PC-3 cell
Human liver tissue lysates
Primary antibody: [TA368410] (ATP5ME Antibody)
at dilution 1/400
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution
Exposure time: 5 seconds

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA368410] (ATP5ME Antibody) at dilution 1/65 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA368410] (ATP5ME Antibody) at dilution 1/65, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA368410] (ATP5ME Antibody) at dilution 1/65 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA368410] (ATP5ME Antibody) at dilution 1/65, treated with synthetic peptide. (Original magnification: ×200)