

### **Product datasheet for TA368584**

# Pancreatic Polypeptide (PPY) Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1000-5000

WB positive control: Rat pancreas tissue lysate

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide of human PPY

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Producted Protein Size: 11 kDa

Gene Name:

**Predicted Protein Size:** 11 kDa

Database Link: Entrez Gene 5539 Human

P01298

pancreatic polypeptide

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

This gene encodes a member of the neuropeptide Y (NPY) family of peptides. The encoded 95 aa preproprotein is synthesized in the pancreatic islets of Langerhans and proteolytically processed to generate two peptide products. These products include the active pancreatic hormone of 36 aa and an icosapeptide of unknown function. This hormone acts as a regulator of pancreatic and gastrointestinal functions and may be important in the regulation of food intake. Plasma level of this hormone has been shown to be reduced in conditions associated with increased food intake and elevated in anorexia nervosa. In addition, infusion of this hormone in obese rodents has shown to decrease weight gain. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.

Synonyms: Obinepitide; PNP; PP

## **Product images:**



Gel: 12%SDS-PAGE Lysate: 40 μg

Lane: Rat pancreas tissue lysate

Primary antibody: TA368584 (PPY Antibody) at

dilution 1/2500

Secondary antibody: Goat anti rabbit IgG at

1/5000 dilution

Exposure time: 3 minutes