

Product datasheet for TA368588S

Junctional Adhesion Molecule C (JAM3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human cerebrum tissue lysate

IHC: 40-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human JAM3

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 35 kDa

Gene Name: junctional adhesion molecule 3

Database Link: Entrez Gene 83700 Human

Q9BX67

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. The protein encoded by this immunoglobulin superfamily gene member is localized in the tight junctions between high endothelial cells. Unlike other proteins in this family, the this protein is unable to adhere to leukocyte cell lines and only forms weak homotypic interactions. The encoded protein is a member of the junctional adhesion molecule protein family and acts as a receptor for another member of this family. A mutation in an intron of this gene is associated with hemorrhagic destruction of the brain, subependymal calcification, and congenital cataracts. Alternative splicing results in multiple transcript variants.

Synonyms:

FLJ14529; JAM-2; JAM-3; JAM-C; JAMC

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human cerebrum tissue lysate


Primary antibody: [TA368588] (JAM3 Antibody) at

dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 3 minutes



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368588] (JAM3 Antibody) at dilution 1/40 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA368588] (JAM3 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)