

Product datasheet for TA368624

KIR2DL5A Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 1000-5000 WB positive control: Human plasma solution IHC: 40-200 Positive control: Human liver cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human KIR2DL5A
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	41 kDa
Gene Name:	killer cell immunoglobulin like receptor, two lg domains and long cytoplasmic tail 5A
Database Link:	<u>Entrez Gene 57292 Human</u> <u>Q8N109</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE KIR2DL5A Rabbit Polyclonal Antibody – TA368624

Background:Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed
by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly
homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb
leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among
haplotypes, although several framework genes are found in all haplotypes (KIR3DL3, KIR3DP1,
KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular
immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S)
cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory
signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR
proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the
TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for
several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to
play an important role in regulation of the immune response.

Synonyms: CD158F; KIR2DL5; KIR2DL5.1; KIR2DL5.3; KIR2DL5B

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: Human plasma solution Primary antibody: TA368624 (KIR2DL5A Antibody) at dilution 1/1000 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution Exposure time: 1 minute

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA368624 (KIR2DL5A Antibody) at dilution 1/40 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA368624 (KIR2DL5A Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US