

# **Product datasheet for TA368630S**

## **SHTN1 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal liver tissue lysate

IHC: 50-100

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human SHTN1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 72 kDa
Gene Name: shootin 1

Database Link: Entrez Gene 57698 Human

A0MZ66

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

Involved in the generation of internal asymmetric signals required for neuronal polarization and neurite outgrowth. Mediates netrin-1-induced F-actin-substrate coupling or 'clutch engagement' within the axon growth cone through activation of CDC42, RAC1 and PAK1-dependent signaling pathway, thereby converting the F-actin retrograde flow into traction forces, concomitantly with filopodium extension and axon outgrowth. Plays a role in cytoskeletal organization by regulating the subcellular localization of phosphoinositide 3-kinase (PI3K) activity at the axonal growth cone. Plays also a role in regenerative neurite outgrowth. In the developing cortex, cooperates with KIF20B to promote both the transition from the multipolar to the bipolar stage and the radial migration of cortical neurons from the ventricular zone toward the superficial layer of the neocortex. Involved in the accumulation of phosphatidylinositol 3,4,5-trisphosphate (PIP3) in the growth cone of primary hippocampal neurons.

### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane: Human fetal liver tissue lysate


Primary antibody: [TA368630] (SHTN1 Antibody)

at dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 5 seconds



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA368630] (SHTN1 Antibody) at dilution 1/60 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA368630] (SHTN1 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)