

# **Product datasheet for TA368648S**

## **CKMT1A Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse heart tissue lysate

IHC: 50-100

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human CKMT1A/CKMT1B

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 47 kDa

**Gene Name:** creatine kinase, mitochondrial 1A

Database Link: Entrez Gene 548596 Human

P12532

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

Mitochondrial creatine (MtCK) kinase is responsible for the transfer of high energy phosphate from mitochondria to the cytosolic carrier, creatine. It belongs to the creatine kinase isoenzyme family. It exists as two isoenzymes, sarcomeric MtCK and ubiquitous MtCK, encoded by separate genes. Mitochondrial creatine kinase occurs in two different oligomeric forms: dimers and octamers, in contrast to the exclusively dimeric cytosolic creatine kinase isoenzymes. Many malignant cancers with poor prognosis have shown overexpression of ubiquitous mitochondrial creatine kinase; this may be related to high energy turnover and failure to eliminate cancer cells via apoptosis. Ubiquitous mitochondrial creatine kinase has 80% homology with the coding exons of sarcomeric mitochondrial creatine kinase. Two genes located near each other on chromosome 15 have been identified which encode identical mitochondrial creatine kinase proteins.

Synonyms:

CKMT; CKMT1; FLJ50967; Mia-CK; U-MtCK; UMTCK

### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane: Mouse heart tissue lysate


Primary antibody: [TA368648] (CKMT1A/CKMT1B

Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA368648] (CKMT1A/CKMT1B Antibody) at dilution 1/40 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA368648] (CKMT1A/CKMT1B Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)