

Product datasheet for TA368677

PD-L1 (CD274) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Raji and A549 cell lysates

IHC: 50-200

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human
Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human CD274

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 33 kDa

Gene Name: CD274 molecule

Database Link: Entrez Gene 29126 Human

Q9NZQ7

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.

Synonyms:

B7-H; B7-H1; B7H1; MGC142294; MGC142296; PD-L1; PDCD1L1; PDCD1LG1; PDL1

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane 1-2: Raji and A549 cell lysates


Primary antibody: TA368677 (CD274 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 10 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA368677 (CD274 Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA368677 (CD274 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)