

Product datasheet for TA368701

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CYP8B1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse heart tissue, Human hepatocellular carcinoma 2 tissue lysates

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human CYP8B1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 58 kDa

Gene Name: cytochrome P450 family 8 subfamily B member 1

Database Link: Entrez Gene 1582 Human

Q9UNU6






Background:

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the conversion of 7 alpha-hydroxy-4-cholesten-3-one into 7-alpha,12-alpha-dihydroxy-4-cholesten-3-one. The balance between these two steroids determines the relative amounts of cholic acid and chenodeoxycholic acid both of which are secreted in the bile and affect the solubility of cholesterol. This gene is unique among the cytochrome P450 genes in that it is intronless.

Synonyms: CP8B; CYP12; CYPVIIIB1; FLJ17826

Product images:



Gel: 8%SDS-PAGE Lysate: 16 μg Lane 1-2: Mouse heart tissue

Human hepatocellular carcinoma 2 tissue lysates


Primary antibody: TA368701 (CYP8B1 Antibody)

at dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA368701 (CYP8B1 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA368701 (CYP8B1 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA368701 (CYP8B1 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA368701 (CYP8B1 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)