

Product datasheet for TA368820S

Histone H1.2 (HIST1H1C) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Jurkat cell lysate

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human H1-2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 21 kDa

Gene Name: histone cluster 1, H1c

Database Link: Entrez Gene 3006 Human

P16403

Background: Histones are basic nuclear proteins responsible for nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H1 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene

cluster on chromosome 6. [provided by RefSeq, Aug 2015]

Synonyms: H1.2; H1F2; MGC3992




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane: Jurkat cell lysate Primary antibody: [TA368820] (H1-2 Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 1 second