

## **Product datasheet for TA368880**

# spindlin 1 (SPIN1) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Jurkat, 293T cell lysates

IHC: 50-200

Positive control: Human colorectal cancer Predicted cell location: Nucleus and Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human SPIN1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 30 kDa
Gene Name: spindlin 1

Database Link: Entrez Gene 10927 Human

Q9Y657

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

Chromatin reader that specifically recognizes and binds histone H3 both trimethylated at 'Lys-4' and asymmetrically dimethylated at 'Arg-8' (H3K4me3 and H3R8me2a) and acts as an activator of Wnt signaling pathway downstream of PRMT2. In case of cancer, promotes cell cancer proliferation via activation of the Wnt signaling pathway (PubMed:24589551). Overexpression induces metaphase arrest and chromosomal instability. Localizes to active rDNA loci and promotes the expression of rRNA genes (PubMed:21960006). May play a role in cell-cycle regulation during the transition from gamete to embryo. Involved in oocyte meiotic resumption, a process that takes place before ovulation to resume meiosis of oocytes blocked in prophase I: may act by regulating maternal transcripts to control meiotic resumption.

**Synonyms:** OCR; SPIN; spindlin; SPNS1


### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: Jurkat 293T cell lysates Primary antibody: TA368880 (SPIN1 Antibody) at dilution 1/600 Secondary antibody: Goat anti rabbit IgG at


Exposure time: 30 seconds

1/5000 dilution




Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA368880 (SPIN1 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA368880 (SPIN1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA368880 (SPIN1 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA368880 (SPIN1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)