

Product datasheet for TA368936

IDH3B Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: IHC, WB Recommended Dilution: WB: 200-1000 WB positive control: Mouse eyes tissue IHC: 25-100 Positive control: Human thyroid cancer Predicted cell location: Cytoplasm **Reactivity:** Human, Mouse, Rat Host: Rabbit Isotype: lgG **Clonality:** Polyclonal Immunogen: Fusion protein of human IDH3B pH7.4 PBS, 0.05% NaN3, 40% Glycerol Formulation: Concentration: lot specific Antigen affinity purification **Purification: Conjugation:** Unconjugated Storage: Store at -20°C. Stability: 1 year **Predicted Protein Size:** 42 kDa Gene Name: isocitrate dehydrogenase 3 (NAD(+)) beta Database Link: Entrez Gene 3420 Human 043837

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE IDH3B Rabbit Polyclonal Antibody – TA368936

Background: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. The protein encoded by this gene is the beta subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase. Three alternatively spliced transcript variants encoding different isoforms have been described for this gene.

Synonyms:

FLJ11043; H-IDHB; MGC903; RP46

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg Lane: Mouse eyes tissue Primary antibody: TA368936 (IDH3B Antibody) at dilution 1/240 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 10 seconds

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA368936 (IDH3B Antibody) at dilution 1/30 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA368936 (IDH3B Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US