

Product datasheet for TA369067S

Cystatin SA (CST2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human liver cancer tissue

IHC: 50-200

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human CST2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 16 kDa
Gene Name: cystatin SA

Database Link: Entrez Gene 1470 Human

P09228

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Background:

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a secreted thiol protease inhibitor found at high levels in saliva, tears and seminal plasma.

Synonyms:

Cystatin-2; Cystatin-S5; MGC71924

Product images:



Gel: 12%SDS-PAGE Lysate: 40 μg

Lane: Human liver cancer tissue


Primary antibody: [TA369067] (CST2 Antibody) at

dilution 1/250

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 minutes



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA369067] (CST2 Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA369067] (CST2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)