

Product datasheet for **TA369250S**

OCIAD1 Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 200-1000 WB positive control: Hela and Jurkat cells, human testis tissue and 231 cells
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human OCIAD1
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	28 kDa
Gene Name:	OCIA domain containing 1
Database Link:	Entrez Gene 54940 Human Q9NX40
Background:	OCIAD1 (OCIA domain containing 1), also known as OCIA (ovarian carcinoma immunoreactive antigen), TPA018 or ASRIJ, is a 245 amino acid endosomal protein that contains one OCIA domain and belongs to the OCIAD1 family. OCIAD1 exists as two alternatively spliced isoforms (termed OCIAD1 isoform A and B) that localize to different tissues; isoform A localizes to prostate, brain, mammary gland, testis, placenta and ovary, whereas isoform B is restricted to the central nervous system. The gene encoding OCIAD1 maps to human chromosome 4, which represents approximately 6% of the human genome, contains nearly 900 genes and is associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.
Synonyms:	Asrij; FLJ20455; MGC111072; OCIA; TPA018



[View online »](#)

Product images:



Gel: 10%SDS-PAGE
Lysate: 40 μ g
Lane 1-4: Hela cells
Jurkat cells
human testis tissue
231 cells
Primary antibody: [TA369250] (OCIAD1 Antibody)
at dilution 1/200
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution
Exposure time: 30 seconds