

Product datasheet for TA369466S

WDR19 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human WDR19

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: WD repeat domain 19

Database Link: Entrez Gene 57728 Human

Q8NEZ3

Background: This gene encodes a member of the WD repeat protein family. WD repeats are minimally

conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp

(GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This protein contains six WD repeats, a clathrin heavy-chain repeat, and three transmembrane domains. This gene is conserved from C. elegans to human. It may participate in androgen-regulated signaling mechanisms or in the vesicular trafficking of androgen-regulated secretory processes. Alternatively spliced transcript variants encoding distinct isoforms have been reported but

the full-length nature of one of these variants has not been defined.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

FLJ23127; KIAA1638; ORF26; Pwdmp

Product images:



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA369466] (WDR19 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA369466] (WDR19 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)