

Product datasheet for TA369484

ZCWCC1 (MORC2) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: 231, 293T, Hela and A172 cell lysates IHC: 25-100 Positive control: Human lung cancer Predicted cell location: Nucleus
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human MORC2
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	118 kDa
Gene Name:	MORC family CW-type zinc finger 2
Database Link:	<u>Entrez Gene 22880 Human</u> <u>Q9Y6X9</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE ZCWCC1 (MORC2) Rabbit Polyclonal Antibody – TA369484


Background:The CW domain is a structural module found in many vertebrate, parasitic and plant
proteins. It consists of a mononuclear four-cysteine zinc-finger domain that plays a role in
DNA binding, chromatin methylation and early embryonic development. ZCWCC1 (zinc finger
CW-type coiled-coil domain protein 1), also known as MORC2 (MORC family CW-type zinc
finger protein 2) or ZCW3, is a 1,032 amino acid protein that contains one CW-type zinc finger
domain. ZCWCC1 is located on chromosome 22 and is ubiquitously expressed with highest
expression in pancreas, smooth muscle and testis. Expression of ZCWCC1 is upregulated in
hypoxia, a pathological condition characterized by an inadequate supply of oxygen in the
blood.

Synonyms: AC004542.C22.1; KIAA0852; ZCW3; ZCWCC1

Product images:




Gel: 6%SDS-PAGE Lysate: 40 µg Lane 1-4: 231 293T Hela and A172 cell lysates Primary antibody: TA369484 (MORC2 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA369484 (MORC2 Antibody) at dilution 1/25 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA369484 (MORC2 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US