

Product datasheet for TA369605

CTRP3 (C1QTNF3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human colorectal cancer tissue

IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Secreted

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human C1QTNF3

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 27 kDa

Gene Name: C1q and tumor necrosis factor related protein 3

Database Link: Entrez Gene 114899 Human

Q9BXI4



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

C1qTNF3 (Complement C1q TNF-related protein 3/CTRP3; also CORS26 and cartonectin) is a 30-32 kDa, secreted member of the C1q and TNF-related protein (CTRP) superfamily of molecules. It is expressed by a wide variety of cells, including smooth muscle cells, fibroblasts, adipocytes, monocytes and proliferating chondrocytes. C1qTNF3 is an antiinflammatory agent that apparently blocks LPS activation of mononuclear cells. It also has marked proliferative activity on diverse cell types such as vascular smooth muscle, chrondrocytes, and endothelium. Finally, C1qTNF3 is known to act on hepatocytes and suppress hepatocyte gluconeogenesis. Mature human C1qTNF3 is 224 amino acids (aa) in length (aa 23-246). It possesses an N-terminal collagen-like domain (aa 51-113) followed by a C-terminal globular region (aa 113-246). C1qTNF3 is monomeric when intracellular, but forms a 90 kDa homotrimer plus higher-order oligomer when secreted. There are at least two potential isoform variants. One is 40-42 kDa, glycosylated, and contains a 73 aa insertion after Glu28, while a second shows concurrent deletions of aa 46-69 and 82-105. The longer 40 kDa isoform is reported to form heterotrimers and oligomers with the standard 30 kDa isoform. This has the effect of protecting the standard isoform from proteolysis. Over aa 24-246, human C1qTNF3 shares 99% aa sequence identity with mouse C1qTNF3.

Synonyms:

2310005P21Rik; C1ATNF3; cartonectin; Corcs; Cors; Cors-26; CORS26; CTRP3; FLJ37576

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg

Lane: Human colorectal cancer tissue

Primary antibody: TA369605 (C1QTNF3 Antibody)


at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA369605 (C1QTNF3 Antibody) at dilution 1/25 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA369605 (C1QTNF3 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA369605 (C1QTNF3 Antibody) at dilution 1/25 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human lung cancer tissue using TA369605 (C1QTNF3 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)