

## **Product datasheet for TA369610S**

## **FKBP6 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human testis tissue

IHC: 25-100

Positive control: Human breast cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Full length fusion protein

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 37 kDa

Gene Name: FK506 binding protein 6

Database Link: Entrez Gene 8468 Human

075344

**Background:** The protein encoded by this gene is a cis-trans peptidyl-prolyl isomerase that may function in

immunoregulation and basic cellular processes involving protein folding and trafficking. This

gene is located in a chromosomal region that is deleted in Williams-Beuren syndrome.

Defects in this gene may cause male infertility. There are multiple pseudogenes for this gene located nearby on chromosome 7. Alternative splicing results in multiple transcript variants.

**Synonyms:** FKBP-36; FKBP36; MGC87179; PPlase; rotamase




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human testis tissue


Primary antibody: [TA369610] (FKBP6 Antibody)

at dilution 1/250


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 seconds



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA369610] (FKBP6 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA369610] (FKBP6 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)