

## Product datasheet for **TA369813**

### ATP5F1C Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: Hela,HEPG2,Jurkat and A549 cell,Human heart tissue lysates IHC: 25-100 Positive control: Human colorectal cancer Predicted cell location: Cytoplasm and Nucleus
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human ATP5F1C
Formulation:	pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	33 kDa
Gene Name:	ATP synthase, H <sup>+</sup> transporting, mitochondrial F1 complex, gamma polypeptide 1
Database Link:	<a href="#">Entrez Gene 509 Human P36542</a>




[View online »](#)

**Background:**

This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. ATP synthase is composed of two linked multi-subunit complexes: the soluble catalytic core, F<sub>1</sub>, and the membrane-spanning component, F<sub>o</sub>, comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel consists of three main subunits (a, b, c). This gene encodes the gamma subunit of the catalytic core. Alternatively spliced transcript variants encoding different isoforms have been identified. This gene also has a pseudogene on chromosome 14. [provided by RefSeq, Jul 2008]

**Synonyms:**

ATP5C; ATP5CL1

**Product images:**

Gel: 8%SDS-PAGE

Lysate: 40 µg

Lane 1-5: HeLa

HEPG2

Jurkat and A549 cell

Human heart tissue lysates


Primary antibody: TA369813 (ATP5F1C Antibody)

at dilution 1/650


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 5 seconds



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA369813 (ATP5F1C Antibody) at dilution 1/35 (Original magnification:  $\times 200$ )



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA369813 (ATP5F1C Antibody) at dilution 1/35, treated with fusion protein. (Original magnification:  $\times 200$ )