

## **Product datasheet for TA369832S**

## i roddet datasneet for 1A303032.

## **BHMT2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Human fetal liver tissue and Human liver tissue lysates

IHC: 30-150

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human BHMT2

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 40 kDa

**Gene Name:** betaine--homocysteine S-methyltransferase 2

Database Link: Entrez Gene 23743 Human

Q9H2M3



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Homocysteine is a sulfur-containing amino acid that plays a crucial role in methylation reactions. Transfer of the methyl group from betaine to homocysteine creates methionine, which donates the methyl group to methylate DNA, proteins, lipids, and other intracellular metabolites. The protein encoded by this gene is one of two methyl transferases that can catalyze the transfer of the methyl group from betaine to homocysteine. Anomalies in homocysteine metabolism have been implicated in disorders ranging from vascular disease to neural tube birth defects such as spina bifida. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Synonyms:

FLJ20001

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane 1-2: Human fetal liver tissue and Human

liver tissue lysates


Primary antibody: [TA369832] (BHMT2 Antibody)

at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 10 seconds



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA369832] (BHMT2 Antibody) at dilution 1/30 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA369832] (BHMT2 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)