

Product datasheet for TA369887

Complement C7 (C7) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse lung tissue lysate

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human C7

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 94 kDa

Gene Name: complement component 7

Database Link: Entrez Gene 730 Human

P10643

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

The complement cascade is a multi-protein system that functions to clear pathogens from an infected host. Part of the innate (unchanging) immune system, the complement cascade consists of proteins and inactive zymogens that are present in blood and are stimulated by one of several triggers. Once stimulated, the cascade relays amplified responses throughout the body, ultimately activating the cell-killing membrane attack complex which can insert itself into the cell membrane and cause the cell to lyse. C7 (complement component 7) is an 843 amino acid secreted protein that participates in the formation of membrane attack complex (MAC), a complex that forms pores in the plasma membrane of target cells for innate and adaptive immune responses. As a membrane anchor, C7 exists as a monomer or dimer and can form multimeric rosettes with C5 β . C7 defects are the cause of component C7 deficiency (C7D), characterized by recurrent bacterial infections caused by Neisseria meningitidis.

Synonyms: C7

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane: Mouse lung tissue lysate

Primary antibody: TA369887 (C7 Antibody) at

dilution 1/800

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 minutes