

Product datasheet for TA369956

CEP57 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: A549 cell lysate

IHC: 25-100

Positive control: Human liver cancer

Predicted cell location: Cytoplasm or Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Full length fusion protein

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 57 kDa

Gene Name: centrosomal protein 57

Database Link: Entrez Gene 9702 Human

<u>Q86XR8</u>

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

This gene encodes a cytoplasmic protein called Translokin. This protein localizes to the centrosome and has a function in microtubular stabilization. The N-terminal half of this protein is required for its centrosome localization and for its multimerization, and the C-terminal half is required for nucleating, bundling and anchoring microtubules to the centrosomes. This protein specifically interacts with fibroblast growth factor 2 (FGF2), sorting nexin 6, Ran-binding protein M and the kinesins KIF3A and KIF3B, and thus mediates the nuclear translocation and mitogenic activity of the FGF2. It also interacts with cyclin D1 and controls nucleocytoplasmic distribution of the cyclin D1 in quiescent cells. This protein is crucial for maintaining correct chromosomal number during cell division. Mutations in this gene cause mosaic variegated aneuploidy syndrome, a rare autosomal recessive disorder. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Synonyms: KIAA0092; PIG8; translokin; TSP57

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: A549 cell lysate

Primary antibody: TA369956 (CEP57 Antibody) at


dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 2 minutes






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA369956 (CEP57 Antibody) at dilution 1/20 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA369956 (CEP57 Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA369956 (CEP57 Antibody) at dilution 1/20 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA369956 (CEP57 Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)