

Product datasheet for TA370174S

RPL3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: 293T, Jurkat and Raji cell lysates

IHC: 30-150

Positive control: Human cervical cancer Predicted cell location: Cytoplasm or Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human RPL3

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 46 kDa

Gene Name: ribosomal protein L3

Database Link: Entrez Gene 6122 Human

P39023

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

Ribosomes, the complexes that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L3P family of ribosomal proteins and it is located in the cytoplasm. The protein can bind to the HIV-1 TAR mRNA, and it has been suggested that the protein contributes to tat-mediated transactivation. This gene is cotranscribed with several small nucleolar RNA genes, which are located in several of this gene's introns. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Synonyms:

MGC104284; TARBP-B

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-3: 293T Jurkat and Raji cell lysates


Primary antibody: [TA370174] (RPL3 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit $\lg G$ at


1/8000 dilution

Exposure time: 5 seconds




Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA370174] (RPL3 Antibody) at dilution 1/20 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA370174] (RPL3 Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA370174] (RPL3 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA370174] (RPL3 Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)