

# **Product datasheet for TA370339S**

# **PSMA3** Rabbit Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela cell, Mouse spleen tissue, Mouse liver tissue, PC3 cell, HL60 cell,

A549 cell, NIH/3T3 cell lysates

IHC: 50-300

Positive control: Human lung cancer

Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse, Rat

Rabbit Host: Isotype: lgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PSMA3

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Store at -20°C. Storage:

Stability: 1 year **Predicted Protein Size:** 28 kDa

Gene Name: proteasome subunit alpha 3 Database Link: Entrez Gene 5684 Human

P25788



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com





### Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Two alternative transcripts encoding different isoforms have been identified.

Synonyms: HC8; MGC12306; MGC32631; PSC3; PSC8

## **Product images:**







Gel: 12%SDS-PAGE
Lysate: 40 µg
Lane 1-7: Hela cell
Mouse spleen tissue
Mouse liver tissue
PC3 cell
HL60 cell
A549 cell
NIH/3T3 cell lysates
Primary antibody: [TA370339] (PSMA3 Antibody)
at dilution 1/350
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution
Exposure time: 10 seconds

Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA370339] (PSMA3 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA370339] (PSMA3 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA370339] (PSMA3 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA370339] (PSMA3 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)