

Product datasheet for TA370530S

ARD1A (NAA10) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Hela and 293T cell lysates

IHC: 50-300

Positive control: Human tonsil

Predicted cell location: Cytoplasm or Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human NAA10

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 26 kDa

Gene Name: N(alpha)-acetyltransferase 10, NatA catalytic subunit

Database Link: Entrez Gene 8260 Human

P41227

Background: N-alpha-acetylation is among the most common post-translational protein modifications in

eukaryotic cells. This process involves the transfer of an acetyl group from acetyl-coenzyme A to the alpha-amino group on a nascent polypeptide and is essential for normal cell function. This gene encodes an N-terminal acetyltransferase that functions as the catalytic subunit of the major amino-terminal acetyltransferase A complex. Mutations in this gene are the cause

of Ogden syndrome. Alternate splicing results in multiple transcript variants.

Synonyms: ARD1; ARD1A; DXS707; NATD; TE2




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-2: Hela and 293T cell lysates


Primary antibody: [TA370530] (NAA10 Antibody)

at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370530] (NAA10 Antibody) at dilution 1/55 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370530] (NAA10 Antibody) at dilution 1/55, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA370530] (NAA10 Antibody) at dilution 1/55 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA370530] (NAA10 Antibody) at dilution 1/55, treated with fusion protein. (Original magnification: ×200)