

Product datasheet for TA370553S

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Rockville, MD 20850, US

CN: techsupport@origene.cn

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

ATP5PD Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse skeletal muscle tissue, Mouse kidney tissue, PC-3, Jurkat, HepG2

and Hela cell lysates

IHC: 50-300

Positive control: Human colorectal cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human ATP5PD

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Predicted Protein Size: 18 kDa

Gene Name: ATP synthase, H+ transporting, mitochondrial Fo complex subunit D

Database Link: Entrez Gene 10476 Human

O75947






Background:

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the d subunit of the Fo complex. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. In addition, three pseudogenes are located on chromosomes 9, 12 and 15.

Synonyms: ATP5JD; ATPQ

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg

Lane 1-6: Mouse skeletal muscle tissue

Mouse kidney tissue

PC-3 Jurkat

HepG2 and Hela cell lysates


Primary antibody: [TA370553] (ATP5PD Antibody)

at dilution 1/300

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 3 seconds




Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA370553] (ATP5PD Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA370553] (ATP5PD Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370553] (ATP5PD Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370553] (ATP5PD Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)