

Product datasheet for TA370580S

EWSR1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: 293T and HepG2 cell lysates

IHC: 50-300

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human EWSR1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 68 kDa

Gene Name: EWS RNA binding protein 1

Database Link: Entrez Gene 2130 Human

Q01844

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

This gene encodes a multifunctional protein that is involved in various cellular processes, including gene expression, cell signaling, and RNA processing and transport. The protein includes an N-terminal transcriptional activation domain and a C-terminal RNA-binding domain. Chromosomal translocations between this gene and various genes encoding transcription factors result in the production of chimeric proteins that are involved in tumorigenesis. These chimeric proteins usually consist of the N-terminal transcriptional activation domain of this protein fused to the C-terminal DNA-binding domain of the transcription factor protein. Mutations in this gene, specifically a t(11;22)(q24;q12) translocation, are known to cause Ewing sarcoma as well as neuroectodermal and various other tumors. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1 and 14.

Synonyms: bK984G1.4; EWS

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg


Lane 1-2: 293T and HepG2 cell lysates Primary antibody: [TA370580] (EWSR1 Antibody)

at dilution 1/1000

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA370580] (EWSR1 Antibody) at dilution 1/70 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA370580] (EWSR1 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA370580] (EWSR1 Antibody) at dilution 1/70 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA370580] (EWSR1 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)