

Product datasheet for TA370603

Froduct datasineet for TAS70003

PDHA1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Mouse brain tissue, HepG2, Hela, A549 cell, Mouse skeletal muscle tissue,

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

NIH/3T3 cell lysates

IHC: 50-300

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PDHA1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 43 kDa

Gene Name: pyruvate dehydrogenase (lipoamide) alpha 1

Database Link: Entrez Gene 5160 Human

P08559






Background:

The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2), and provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle. The PDH complex is composed of multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and lipoamide dehydrogenase (E3). The E1 enzyme is a heterotetramer of two alpha and two beta subunits. This gene encodes the E1 alpha 1 subunit containing the E1 active site, and plays a key role in the function of the PDH complex. Mutations in this gene are associated with pyruvate dehydrogenase E1-alpha deficiency and X-linked Leigh syndrome. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Synonyms: PDHA; PDHCE1A; PHE1A


Product images:




Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-6: Mouse brain tissue
HepG2
Hela
A549 cell
Mouse skeletal muscle tissue
NIH/3T3 cell lysates
Primary antibody: TA370603 (PDHA1 Antibody) at dilution 1/1000
Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution
Exposure time: 1 second

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA370603 (PDHA1 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA370603 (PDHA1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA370603 (PDHA1 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA370603 (PDHA1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)