

## **Product datasheet for TA370627S**

## **KIR3DL1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-300

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human KIR3DL1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1

**Database Link:** Entrez Gene 3811 Human

P43629



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




## Background:

Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several framework genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response.

Synonyms:


AMB11; CD158E; CD158E1; CD158e1/2; CD158e2; cl-2; cl-11; KIR; KIR3DS1; MGC119726; MGC119728; MGC126589; MGC126591; NK-receptor; NKAT-3; NKB1; NKB1B

## **Product images:**




Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370627] (KIR3DL1 Antibody) at dilution 1/95 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370627] (KIR3DL1 Antibody) at dilution 1/95, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA370627] (KIR3DL1 Antibody) at dilution 1/95 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA370627] (KIR3DL1 Antibody) at dilution 1/95, treated with fusion protein. (Original magnification: ×200)