

# **Product datasheet for TA370780**

## **CCDC134 Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Human fetal liver tissue, 293T cell lysates

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human CCDC134

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 27 kDa

**Gene Name:** coiled-coil domain containing 134

Database Link: Entrez Gene 79879 Human

Q9H6E4



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

In extracellular secreted form, promotes proliferation and activation of CD8+ T cells, suggesting a cytokine-like function (PubMed:25125657). Enhances cytotoxic anti-tumor activity of CD8+ T cells (PubMed:25125657). May inhibit ERK and JNK signaling activity (PubMed:18087676, PubMed:23070808). May suppress cell migration and invasion activity, via its effects on ERK and JNK signaling (PubMed:23070808). In the nucleus, enhances stability of the PCAF histone acetyltransferase (HAT) complex member TADA2A and thus promotes PCAF-mediated H3K14 and H4K8 HAT activity. May inhibit TADA2A-mediated TP53/p53 'Lys-321' acetylation, leading to reduced TP53 stability and transcriptional activity. May also promote TADA2A-mediated XRCC6 acetylation thus facilitating cell apoptosis in response to DNA damage.

Synonyms:

dJ821D11.3; FLJ22349; MGC21013

### **Product images:**



Gel: 12%SDS-PAGE Lysate: 40 µg

Lane 1-2: Human fetal liver tissue

293T cell lysates


Primary antibody: TA370780 (CCDC134 Antibody)

at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA370780 (CCDC134 Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA370780 (CCDC134 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)