

## **Product datasheet for TA370862S**

# **GRIK2 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: LOVO, Jurkat and 293T cell lysates

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human GRIK2

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 103 kDa

**Gene Name:** glutamate ionotropic receptor kainate type subunit 2

Database Link: Entrez Gene 2898 Human

Q13002



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. This gene product belongs to the kainate family of glutamate receptors, which are composed of four subunits and function as ligand-activated ion channels. The subunit encoded by this gene is subject to RNA editing at multiple sites within the first and second transmembrane domains, which is thought to alter the structure and function of the receptor complex. Alternatively spliced transcript variants encoding different isoforms have also been described for this gene. Mutations in this gene have been associated with autosomal recessive cognitive disability.

Synonyms:


bA487F5.1; EAA4; GLR6; GLUK6; GluR-6; GLUR6; MGC74427; MRT6; OTTHUMP00000017950

### **Product images:**




Gel: 6%SDS-PAGE
Lysate: 40 µg
Lane 1-3: LOVO
Jurkat and 293T cell lysates
Primary antibody: [TA370862] (GRIK2 Antibody) at dilution 1/700
Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA370862] (GRIK2 Antibody) at dilution 1/100 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA370862] (GRIK2 Antibody) at dilution 1/100, treated with fusion protein. (Original magnification: ×200)