

## **Product datasheet for TA370881S**

# **LYAR Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: LOVO cell lysate

IHC: 50-200

Positive control: Human gastric cancer

Predicted cell location: Nucleus

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human LYAR

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 44 kDa

Gene Name: Ly1 antibody reactive

Database Link: Entrez Gene 55646 Human

Q9NX58

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

Plays a role in the maintenance of the appropriate processing of 47S/45S pre-rRNA to 32S/30S pre-rRNAs and their subsequent processing to produce 18S and 28S rRNAs (PubMed:24495227). Also acts at the level of transcription regulation. Along with PRMT5, binds the gamma-globin (HBG1/HBG2) promoter and represses its expression (PubMed:25092918). In neuroblastoma cells, may also repress the expression of oxidative stress genes, including CHAC1, HMOX1, SLC7A11, ULBP1 and SNORD41 that encodes a small nucleolar RNA (PubMed:28686580). Preferentially binds to a DNA motif containing 5'-GGTTAT-3' (PubMed:25092918). Stimulates phagocytosis of photoreceptor outer segments by retinal pigment epithelial cells (By similarity). Prevents nucleolin/NCL self-cleavage, maintaining a normal steady-state level of NCL protein in undifferentiated embryonic stem cells (ESCs), which in turn is essential for ESC self-renewal (By similarity).

Synonyms:

FLJ20425; ZLYAR

### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: LOVO cell lysate


Primary antibody: [TA370881] (LYAR Antibody) at

dilution 1/1000

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA370881] (LYAR Antibody) at dilution 1/140 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA370881] (LYAR Antibody) at dilution 1/140, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370881] (LYAR Antibody) at dilution 1/140 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA370881] (LYAR Antibody) at dilution 1/140, treated with fusion protein. (Original magnification: ×200)