

# Product datasheet for TA370893

# **PHAX Rabbit Polyclonal Antibody**

# **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	IHC: 200-300 Positive control: Human esophagus cancer Predicted cell location: Nucleus
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human PHAX
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Gene Name:	phosphorylated adaptor for RNA export
Database Link:	<u>Entrez Gene 51808 Human</u> <u>Q9H814</u>

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


## **GRIGENE** PHAX Rabbit Polyclonal Antibody – TA370893

#### Background:


A phosphoprotein adapter involved in the XPO1-mediated U snRNA export from the nucleus. Bridge components required for U snRNA export, the cap binding complex (CBC)-bound snRNA on the one hand and the GTPase Ran in its active GTP-bound form together with the export receptor XPO1 on the other. Its phosphorylation in the nucleus is required for U snRNA export complex assembly and export, while its dephosphorylation in the cytoplasm causes export complex disassembly. It is recycled back to the nucleus via the importin alpha/beta heterodimeric import receptor. The directionality of nuclear export is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. Its compartmentalized phosphorylation cycle may also contribute to the directionality of export. Binds strongly to m7G-capped U1 and U5 small nuclear RNAs (snRNAs) in a sequence-unspecific manner and phosphorylation-independent manner (By similarity). Plays also a role in the biogenesis of U3 small nucleolar RNA (snoRNA). Involved in the U3 snoRNA transport from nucleoplasm to Cajal bodies. Binds strongly to m7G-capped U3, U8 and U13 precursor snoRNAs and weakly to trimethylated (TMG)-capped U3, U8 and U13 snoRNAs. Binds also to telomerase RNA.

Synonyms: FLJ13193; RNUXA

### **Product images:**



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA370893 (PHAX Antibody) at dilution 1/180 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA370893 (PHAX Antibody) at dilution 1/180, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA370893 (PHAX Antibody) at dilution 1/180 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA370893 (PHAX Antibody) at dilution 1/180, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US