

## **Product datasheet for TA371373S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **HNRPH1 (HNRNPH1) Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: LoVo and A375 cells, mouse brain tissue, HepG2, A431 and Hela cells

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human HNRNPH1Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 49 kDa

Gene Name: heterogeneous nuclear ribonucleoprotein H1 (H)

Database Link: Entrez Gene 3187 Human

P31943

**Background:** This gene encodes a member of a subfamily of ubiquitously expressed heterogeneous

nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins that complex with heterogeneous nuclear RNA. These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some may shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has three repeats of quasi-RRM domains that bind to RNA and is very similar to the family member HNRPF. This gene may be

associated with hereditary lymphedema type I.






Synonyms:

DKFZp686A15170; hnRNPH; HNRPH; HNRPH1; OTTHUMP00000161513

# **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-6: LoVo cells A375 cells mouse brain tissue HepG2 cells A431 cells Hela cells

Primary antibody: [TA371373] (HNRNPH1 Antibody) at dilution 1/100

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 40 seconds