

# **Product datasheet for TA371460S**

## **ALIX (PDCD6IP) Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: Hela cell lysate

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human PDCD6IPFormulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 96 kDa

**Gene Name:** programmed cell death 6 interacting protein

Database Link: Entrez Gene 10015 Human

Q8WUM4

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

This gene encodes a protein that functions within the ESCRT pathway in the abscission stage of cytokinesis, in intralumenal endosomal vesicle formation, and in enveloped virus budding. Studies using mouse cells have shown that overexpression of this protein can block apoptosis. In addition, the product of this gene binds to the product of the PDCD6 gene, a protein required for apoptosis, in a calcium-dependent manner. This gene product also binds to endophilins, proteins that regulate membrane shape during endocytosis. Overexpression of this gene product and endophilins results in cytoplasmic vacuolization, which may be partly responsible for the protection against cell death. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene. Related pseudogenes have been identified on chromosome 15.

Synonyms:

AIP1; Alix; DRIP4; Hp95; KIAA1375; MGC17003

## **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 µg Lane: Hela cell lysate

Primary antibody: [TA371460] (PDCD6IP

Antibody) at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 seconds